

THREE NEW *DROSOPHILA* SPECIES (DIPTERA: DROSOPHILIDAE) FROM BRITISH COLUMBIA, HAWAII, AND THE CANARY ISLANDS^{1,2}

Haruo Takada, Jong S. Yoon³

ABSTRACT: *Drosophila canadiana* n. sp. from Smithers, British Columbia, Canada, *D. canaryana* n. sp. from Tenerife, Canary Islands, and *D. soonae* n. sp. from Kona, Island of Hawaii, are described. Their relationships to other species in the *virilis* species group, the *obscura* species group, and the modified mouthparts species group, respectively, are discussed.

The National *Drosophila* Species Resource Center (NDSRC) at Bowling Green State University, Ohio, maintains a collection of living cultures of *Drosophila*, totaling more than 350 named species and several undescribed species. This is the largest collection of living eukaryotic organisms ever assembled whose evolutionary relationships and genetic biology have been studied extensively.

In order to maintain the quality of stock and to provide better services for the scientific community, it is necessary to verify the authenticity of the stock, and to identify unnamed species. During the course of this study, three of these unnamed species are described: the first species is from British Columbia, the second is from the Canary Islands, and the third is from the Island of Hawaii.

Drosophila canadiana n. sp.

(Figs. 1-2)

Diagnosis: Distinguished from other members of the *virilis* species group by cylindrical shaped spermatheca, phallosomal index of 2.3-2.5, and evenly scattered dorsal hairs on the aedeagus.

Description:

External characteristics of adults: Male and female. Arista with usually 7 branches including a terminal fork. Antennae brown, third joint darker. Frons dark reddish brown, ocellar triangle dark brown. Anterior reclinate orbital bristle less than 1/2 length of pro-

¹Received April 25, 1988. Accepted March 4, 1989.

²This work was supported in part by NSF Grant BSR-8510104.

³Department of Biological Sciences, Bowling Green State University, Bowling Green, Ohio 43403.




Fig. 2. *Drosophila canadiana* n. sp. A. phallic organs of male; left half in dorsal aspect and right half in ventral aspect B. peripheral phallic organs of male; right half in posterior aspect C. bridge connecting claspers.

since 1961. This strain exhibited distinctive differences in karyotype and longevity (Durbin and Yoon, 1987, and unpublished data).

D. canadiana belongs to the *virilis* species group of the subgenus *Drosophila*. It is closely related to but clearly differs from the European *D. littoralis* Meigen 1830. *D. canadiana* has a cylindrical shaped spermatheca, phallosomal index of 2.3 - 2.5, and evenly scattered dorsal hairs on the aedeagus, while *D. littoralis* from Switzerland has a hemispherical spermatheca, phallosomal index of 1.4 - 1.8, and aedeagal hairs on only the dorsal half.

Drosophila canaryana n. sp.

(Figs. 3-5)

Diagnosis: Distinguished from other members of the *obscura* species group by male sex combs with a row of long black teeth on each basal and second tarsal segment of foreleg.

Description:

External characteristics of adults: Male. Body blackish brown. Eye dark red with dark pile. Arista with 3 dorsal and 2 ventral branches in addition to terminal fork. Antennae brown, third joint blackish brown, pollinose. Clypeus and carina pale brown, front dark brown. Ocellus pale yellowish brown. Two prominent orals. Middle orbital about 1/2 length anterior proclinate, 1/3 that of posterior reclinate. Palpus with single prominent bristle. Carina narrow above, gradually widening below, somewhat rounded, not sulcate. Cheeks grayish yellow, narrow, their width less than almost 1/5 greatest diameter of the eyes.

Mesonotum and scutellum dark brown. Acrostichal hairs in 8 rows, anterior scutellars convergent. Anterior dorsocentrals 1/2 length posterior dorsocentrals. Pleura dark brown. Anterior sternopleural 1/2 length posterior; middle one undeveloped.

Legs pale grayish yellow. Fore femur swollen, with a row of bristles on posterior surface, and inner surface of posterior end clouded. Basal tarsal segment of fore leg with row of long black teeth, 26 in number, the row placed slightly obliquely on the segment. Second tarsal joint also with large prominent sex-comb, consisting of a row of long black teeth slightly curved to apex, 20-25 in number, placed obliquely, not parallel, to axis. Position and size of these combs are shown in Figure 1 d and e.

Abdominal tergites brownish black, slightly shining. Haltere white. First tergite paler than others.

Wing hyaline; two prominent bristles at distal costal break. Third costal section with heavy bristles on basal 1/3. Coastal index 2.9; 4th vein index 1.7; 4C index 0.9 and 5x index 2.0.

Length of body 2.2 mm. Wings 2.2 mm.

Female. As above, except that no tarsal combs are present. Spermathecae small spherical bodies, dark brown in color and chitinated.

Male genitalia: Periphallallic organs. Middle and lower portions of genital arch with 28 or more bristles, upper portion with 7 bristles; toe elongate and swollen. Primary clasper without prominent process; 8 primary teeth, 9 spines on posterior tip of anal plate in two rows.

Phallic organs. Hypandrium semi-elliptical in shape with a short pair of paramedian spines. Anterior parameres as long as posterior parameres and with 8 sensilla. Aedeagus bifurcate and rather longer than both parameres. Phallosomal index 0.6.

Eggs: Two filaments, with distal ends expanded.

Chromosomes: $2n = 12$, 5R1D. Karyotype observed in brain ganglion of third instar larvae of male.

Types: Holotype male. The original specimen collected in December, 1971, at Tenerife, Canary Islands; Stock No. 1410-1211 (3265.1) of the NDSRC. Allotype female and paratype males and females are same as above. Type specimens have been deposited at the AMNH.

Remarks: On the basis of the external morphology this species appears to be closely related to *Drosophila subobscura* Collin, 1936 (see Pomini, 1940), subgenus *Sophophora*, Genus *Drosophila*, but can be separated from all known members of the *obscura* species group by the phallic organs and sex-combs of the male.





Fig. 3 *Drosophila canaryana* n. sp. A. ventral aspect of phallic organs B. left half of external male genitalia C. the bridge connected claspers D. fore basal tarsus and 2nd tarsus of male E. fore leg of male F. internal genitalia of male G. sperm pump.



4

Fig. 4. *Drosophila canaryana* n. sp. A. egg-guide B. spermatheca C. egg.

***Drosophila soonae* n. sp.**


(Figs. 6-7)

Diagnosis: Distinguished from other species of Hawaiian *Drosophila* by modified mouthparts: each labellum with a series of long, curved, bristle-like processes extending from the apex.

Description:

External characteristics of adults: Male. Body tannish yellow, eyes scarlet in live specimens. Arista with 6 dorsal and 2 ventral branches in addition to the fork. Third antennal segment darker. Frons tan, orbits and ocellar triangle paler. Carina, cheeks and clypeus tan; carina flat. Ocelli tannish yellow. Proclinate orbital 5/6 length posterior reclinate; anterior reclinate thin, 1/3 length proclinate; proclinate orbital situated at middle of the front and near anterior reclinate. Oral bristles are made up of a row of short black bristles of about equal length. Palpi with a long apical bristle and with densely blackish setae around apical one third. Each labellum with a series of long, curved, bristle-like processes extending from the apex.

Mesonotum tannish yellow. Acrostichal hairs in 8 somewhat irregular rows. Posterior dorsocentral bristles long, anterior ones small and thin. Ventral surface of sternopleurite with 14 long, curved, black bristles. Haltere yellow. Lateral surface of second tergite with numerous upright bristles on the posterior margin, each tergite with an apical brownish band and widely interrupted at middle on sixth tergite. Fourth, fifth and sixth sternites with marginal bristles. Middle sternopleural small. Sterno-index 0.6.



5

Fig. 5. *Drosophila. canaryana* n. sp. karyotype from male brain ganglion of the third instar larvae.

Legs yellow; only fifth tarsal segments dark brown. The tarsi with two rows of long hairs; one outside, the other one inside on the anterior surface.

Wing hyaline, veins pale brown; costal index 5.0; 4V index about 1.2; 4C index 0.5; 5x index 1.0. Third costal section with small black bristles on its basal 2/3.

Length of body about 4 mm in male, wings 4 mm.

Female. Labellum without bristle-like process. Palpi not as shaggy as in male. Tarsus without long hairs. Seventh sternite with marginal bristles. Second oral bristle half the length of first one, prominent. Anterior dorsocentral bristles present, 3/4 of length of posterior dorsocentral bristles. Length body about 3.5-4.0 mm in female, wings, 3.5-4 mm.




Fig. 6. *Drosophila soonae* n. sp. A. ventral aspect of phallic organs B. lateral aspect of phallic organs without hypandrium C. left half of external male genitalia D. proboscis of male (lateral aspect) E. fore leg of male F. egg-guide G. spermatheca H. egg.

**7**

Fig. 7. *Drosophila soonae* n. sp. karyotype from male brain ganglion of the third instar larvae.

Male genitalia: Periphallalic organs. Genital arch with about 10 bristles including two long bristles on lower portion; toe rounded. Clasper with about 11 black primary teeth. The genitalia resemble that of *Drosophila mimica* Hardy as mentioned by Takada (1966).

Phallic organs. Aedeagus apically swollen. Anterior parameres long-elliptical in shape, the apices at slightly above level as the level of insertion of the paramedian spines of hypandrium. Phallosomal index about 3.0.

Egg: Four filaments, apical two rather shorter than posterior two.

Sexual behavior: The male goes to the rear of the female and assumes head-under-wing posture. The male then extends his proboscis and licks the female's egg-guide area. The aggressor positions himself so that the longitudinal axis of his body is at right angles to the other individual. Sometimes the abdomen is bent laterally. The aggressor then rushes sideways with a crab-like motion.

Chromosomes: $2n=12$, 5R1D. One pair of autosomes were double length. Karyotype observed in brain ganglion of third instar larvae of male.

Types: Holotype male. Original specimen collected July 13, 1965, at Kealahou Ranch, Pāhala (3,000 ft.), Kona, Island of Hawaii, by Dr. H.L. Carson, University of Hawaii, on banana baits. Stock No. 15290-2591 (C129.2) of the National *Drosophila* Species Resource Center. Allotype and paratypes are the same as the above. Type specimens have been deposited at the AMNH.

Remarks: This species is closely related to *Drosophila mimica* Hardy, 1965. It belongs to the *mimica* subgroup, subgenus *Drosophila*, Genus *Drosophila*, the so-called modified mouthparts species group (Spieth, 1966). Our stock of this species was originally kept by Mrs. Kyung Soon A. Yoon, during her twenty years as a curator of the National *Drosophila* Species Resource Center. We are naming this species in her honor.

ACKNOWLEDGMENTS

We are grateful to Professor Robert C. Graves and Mr. G. Kent Vermilion of Bowling Green State University for their helpful comments on the study and for critically reviewing this manuscript. Also, the authors thank Mrs. Kyung-Soon Ahn Yoon for her careful maintenance of these stocks for 20 years as a curator of the National *Drosophila* Species Resource Center.

LITERATURE CITED

- Durbin, E.J. and J.S. Yoon. 1987. Longevity in the *Drosophila virilis* species group. II. The *D. montana* phylad. Ohio J. Sci. 87, 90-92.
- Hardy, D.E. 1965. Insects of Hawaii, Vol. 12, Diptera: Cyclorrhapha. Univ. of Hawaii Press, 1-184.
- Pomini, F.P. 1940. Contributi alla conoscenza delle *Drosophila* europee. Boll. Ist. Entomol. Univ. Bologna 12: 10-164.
- Spieth, H.T. 1966. Courtship Behavior of Endemic Hawaiian *Drosophila*. Studies in Genetics III. Univ. Tex. Publ. 6615: 245-313.
- Takada, H. 1966. Male genitalia of some Hawaiian Drosophilidae. Studies in Genetics III. Univ. Tex. Publ. 6615: 315-333.

PALAEODIPTERON WALKERI
(DIPTERA: NYMPHOMYIIDAE)
IN THE ADIRONDACK MOUNTAINS,
NEW YORK¹

Michael E. Smith², Barbara J. Wyskowski³, Charles T. Driscoll³,
Carol M. Brooks³, Christina C. Cosentini³

ABSTRACT: The nymphomyiid *Palaeodipteron walkeri* (Diptera) is reported for the first time from New York State. Larvae were collected from two second-order streams, one of which was acidic (Acid Neutralizing Capacity ≤ 0) with coincident low pH and basic cation concentrations, and high aluminum concentrations. Collection of *P. walkeri* in an acidic stream suggests that this species may be relatively insensitive to potential anthropogenic stream acidification.

The Nymphomyiidae is a small, archaic dipteran family consisting of only four species. One of these species, *Palaeodipteron walkeri* Ide, has a Nearctic distribution; specimens have been collected from New Brunswick (Ide 1965; Kevan and Cutten-Ali-Kahn 1975), southwestern Quebec (Cutten and Kevan 1970), northern Quebec (Back and Woods 1979) and in the United States from Maine (Mingo and Gibbs 1976) and Pennsylvania (Adler *et al.* 1985). As part of a study to determine relationships between benthic invertebrate communities and stream acidity, we collected several larval *P. walkeri* from two low-order woodland streams in the western Adirondack Mountains, New York.

During routine monthly sampling of stream benthic invertebrates using a modified Hess sampler (250 μ m mesh size), two larvae were collected in Pancake-Hall Creek Tributary (43° 50'N 74° 52'W) and five larvae were collected in Beaver Brook (43° 43'N 74° 22'W) on 13 January 1986. These samples were collected in second-order riffle areas of both streams. Substrate at each site was similar, consisting of cobbles overlying gravel and sand. Both streams were well shaded. Vegetation along Beaver Brook and Pancake-Hall Creek Tributary was primarily deciduous stands (*Acer saccharum*, *Fagus grandifolia*, *Betula alleghaniensis*) with small coniferous stands (*Picea rubens*, *Abies balsamea*, *Tsuga canadensis*) interspersed. The understory consisted of sparse forbs and grasses. All specimens of *P. walkeri* are currently housed in the invertebrate collection at Valdosta State College.

Physical and chemical characteristics differed between the two streams (Table 1). Beaver Brook was 2-4 m wide, 15-30 cm deep and non-

¹Received November 21, 1988. Accepted January 18, 1989.

²Department of Biology, Valdosta State College, Valdosta, GA 31698.

³Department of Civil Engineering, Syracuse University, Syracuse, NY 13244-1190.

acidic ($\text{ANC} > 0$) with coincident higher pH, C_B (basic cation) concentrations and lower metal concentrations than those found in Pancake-Hall Creek Tributary. Pancake-Hall Creek Tributary was 1-1.5 m wide, 2-20 cm deep and was acidic ($\text{ANC} \leq 0$) with low pH and C_B concentrations. Metal concentrations were elevated at Pancake-Hall Creek Tributary, especially labile monomeric Al that is toxic to fish (Driscoll *et*

TABLE 1. Yearly means (ranges) of physical and chemical stream parameters in Pancake-Hall Creek Tributary (PAT) and Beaver Brook (BE) from January 1985 - January 1986. Samples were collected monthly, $n = 13$.

PARAMETER	PAT	BE
Temperature ($^{\circ}\text{C}$)	5.7 (0.0 - 14.0)	6.5 (0.0 - 18.0)
Velocity (m/sec)	0.21 (0.13 - 0.29)	0.37 (0.23 - 0.51)
Dissolved Oxygen (mg/l)	10.7 (10.0 - 12.0)	11.2 (10.0 - 13.2)
pH	4.88 (4.53 - 5.80)	6.22 (5.45 - 6.95)
Acid Neutralizing Capacity (ANC) ($\mu\text{eq/l}$)	-5.9 (-18.3 - 21.5)	56.8 (6.2 - 113.7)
Basic Cations (C_B) ($\mu\text{eq/l}$)	149.1 (99.0 - 201.9)	228.3 (152.9 - 286.9)
Dissolved Organic Carbon (DOC) (mg C/l)	3.6 (2.4 - 4.7)	2.1 (1.5 - 2.9)
Total Monomeric Al (mg/l)	0.52 (0.09 - 0.91)	0.04 (0.01 - 0.18)
Labile Monomeric Al (mg/l)	0.38 (0.04 - 0.63)	0.01 (0.0 - 0.16)
Non-labile Monomeric Al (mg/l)	0.14 (0.05 - 0.28)	0.05 (0.01 - 0.34)
Fe (mg/l)	0.14 (0.03 - 0.71)	0.03 (0.0 - 0.15)
Mn (mg/l)	0.08 (0.06 - 0.19)	0.01 (0.0 - 0.03)
Zn (mg/l)	0.05 (0.0 - 0.09)	0.01 (0.0 - 0.03)